Brain Research, 575 (1992) 359–361
Elsevier

Author Index

Agnati, L.F., see Pich, E.M., 265
Akintunde, A. and Buxton, D.F., Differential sites of origin and collateralization of corticospinal neurons in the rat: a multiple fluorescent retrograde tracer study, 86
Al-Tikriti, M.S., Roth, R.H., Kessler, R.M. and Innis, R.B., Autoradiographic localization of dopamine D_1 and D_2 receptors in rat cerebral cortex following unilateral neurotoxic lesions, 39
Allen, D., see Bernstein, I.L., 132
Asmus, S.E., Kincaid, A.E. and Newman, S.W., A species-specific population of tyrosine hydroxylase-immunoreactive neurons in the medial amygdaloid nucleus of the Syrian hamster, 199

Badoer, E., see McAllen, R.M., 329
Baux, G., Fossier, P. and Tauc, L., G proteins are involved in the regulation of transmitter release at an *Aplysia* cholinergic synapse, 155
Belai, A. and Burnstock, G., Restoration of diabetes-induced changes in enteric nerves by phorbol 12,13-dibutyrate, 356
Belkin, M., see Lavie, V., 1
Ben-Bassat, S., see Lavie, V., 1
Ben-Eliyahu, S., Marek, P., Vaccarino, A.L., Mogil, J.S., Sternberg, W.F. and Liebeskind, J.C., The NMDA receptor antagonist MK-801 prevents long-lasting non-associative morphine tolerance in the rat, 304
Bernstein, I.L., Chavez, M., Allen, D. and Taylor, E.M., Area postrema mediation of physiological and behavioral effects of lithium chloride in the rat, 132
Beyer, C., see Martinez-Gomez, M., 279
Biagini, G., see Pich, E.M., 265
Bissette, G., see Myers, B., 325
Burnstock, G., see Belai, A., 356
Bussières, N. and Dubuc, R., Phasic modulation of vestibulospinal neuron activity during fictive locomotion in lampreys, 174
Buxton, D.F., see Akintunde, A., 86

Calcutt, N.A., Muir, D., Powell, H.C. and Mizisin, A.P., Reduced ciliary neuronotrophic factor-like activity in nerves from diabetic or galactose-fed rats, 320
Cameron, A.A., Pover, C.M., Willis, W.D. and Coggeshall, R.E., Evidence that fine primary afferent axons innervate a wider territory in the superficial dorsal horn following peripheral axotomy, 151
Cano, J., see Vizuete, M.L., 231
Carlton, S.M., see Steinman, J.L., 25
Casanova, M.F., see Laruelle, M., 47
Chavez, M., see Bernstein, I.L., 132
Chen, J., Rao, Z.-r. and Shi, J.-w., The sites of origin of α-melanocyte-stimulating hormone-containing axonal components in the lateral area of the midbrain periaqueductal gray of the rat, 333
Chiappinelli, V.A., see Fletcher, G.H., 103
Chirino, R., see Martinez-Gomez, M., 279
Choi, H.K., see Won, L., 6
Claude, P., see Herman, M.A.R., 257
Coggeshall, R.E., see Cameron, A.A., 151
Cole, K.S. and Robertson, D., Early efferent innervation of the developing rat cochlea studied with a carbocyanine dye, 223
Coleman, M.H., Yamaguchi, S.-i. and Rogawski, M.A., Protection against dendrotoxin-induced clonic seizures in mice by anticonvulsant drugs, 138
Crain, S.M. and Shen, K.-F., After chronic opioid exposure sensory neurons become supersensitive to the excitatory effects of opioid agonists and antagonists as occurs after acute elevation of GM1 ganglioside, 13

Deas, J., Silver, I.A. and Erecińska, M., Effect of inhibitors of *N*-linked oligosaccharide processing on the high-affinity transport of D-aspartate by C6 glioma cells, 251
Dietrich, W.D., see Markgraf, C.G., 238
DiFiglia, M., see Freese, A., 351
Dohanics, J., Hoffman, G.E., Smith, M.S. and Verbalis, J.G., Functional neurolobectomy induced by controlled compression of the pituitary stalk, 215
Dreifuss, J.J., see Dubois-Dauphin, M., 124
Dubois-Dauphin, M., Raggenbass, M., Widmer, H., Tribollet, E. and Dreifuss, J.J., Morphological and electrophysiological evidence for postsynaptic localization of functional oxytocin receptors in the rat dorsal motor nucleus of the vagus nerve, 124
Dubuc, R., see Bussières, N., 174

Erecińska, M., see Deas, J., 251
Eyzaguirre, C., see Pang, L., 167

Farooqui, S.M., see Sakata, M., 309
Ferraguti, F., see Pich, E.M., 265
Finklestein, S.P., see Freese, A., 351
Fletcher, G.H. and Chiappinelli, V.A., An inward rectifier is present in presynaptic nerve terminals in the chick ciliary ganglion, 103
Fossier, P., see Baux, G., 155
Freese, A., Finklestein, S.P. and DiFiglia, M., Basic fibroblast growth factor protects striatal neurons in vitro from NMDA-receptor mediated excitotoxicity, 351
Fujino, M., see Masuo, Y., 113
Fuxe, K., see Pich, E.M., 265

Gauer, F., Masson-Pévet, M. and Pévet, P., Pinealectomy and constant illumination increase the density of melatonin binding sites in the pars tuberalis of rodents, 32
Ginsberg, M.D., see Markgraf, C.G., 238
Glick, S.D., see Maisonneuve, I.M., 69
Green, E.J., see Markgraf, C.G., 238
Grundmann, K., see Newman, G.C., 159

Halvorson, I., see Hugie, T., 57
Haycock, D.A., see Jansen, E.M., 93
Heller, A., see Won, L., 6
Heller, B., see Won, L., 6
Henshel, D.S. and Miller, R.F., Catecholamine effects on dissociated tiger salamander Muller (glial) cells, 208
Herman, M.A.R., Schulz, C.A. and Claude, P., Early and late effects of NGF may be mediated by different pathways in transdifferentiating chromaffin cells, 257
Hoffman, G.E., see Dohanics, J., 215
Hoffmann, P.C., see Won, L., 6
Hospod, F.E., see Newman, G.C., 159
Hugie, T., Halvorson, I. and Thornhill, J., Brown adipose tissue temperature responses following electrical stimulation of ventromedial hypothalamic and lateral preoptic areas or after norepinephrine infusion to Long Evans or Sprague–Dawley rats, 57
Hurwitz, B.E., see Markgraf, C.G., 238

Ikeda, H., Markey, C.J. and Markey, S.P., Search for neurotoxins structurally related to 1-methyl-4-phenylpyridine (MPP^+) in the pathogenesis of Parkinson's disease, 285
Innis, R.B., see Al-Tikriti, M.S., 39
Ishikawa, K., see Tokunaga, K., 164

Jansen, E.M., Haycock, D.A., Ward, S.J. and Seybold, V.S., Distribution of cannabinoid receptors in rat brain determined with aminoalkylindoles, 93
Jaskiw, G.E., see Laruelle, M., 47
Jen, P.H.-S., see Kamada, T., 187
Jiao, S., Schultz, E. and Wolff, J.A., Intracerebral transplants of primary muscle cells: a potential 'platform' for transgene expression in the brain, 143

Kamada, T., Wu, M. and Jen, P.H.-S., Auditory response properties and spatial response areas of single neurons in the pontine nuclei of the big brown bat, *Eptesicus fuscus*, 187
Kauppila, T., Polyamines enhance recovery after sciatic nerve trauma in the rat, 299
Keller Jr., R.W., see Maisonneuve, I.M., 69
Kessler, R.M., see Al-Tikriti, M.S., 39
Kimura, M., see Nojima, H., 325
Kimura, A. and Tamai, Y., Sensory re-

sponse of cortical neurons in the anterior ectosylvian sulcus, including the area evoking eye movement, 181
Kimura, I., see Nojima, H., 337
Kimura, M., see Nojima, H., 337
Kincaid, A.E., see Asmus, S.E., 199
Kleinman, J.E., see Laruelle, M., 47
Kodama, R.T., see Smith, K.J., 79
Kolachana, B., see Laruelle, M., 47
Komisaruk, B.R., see Martinez-Gomez, M., 279
Kontur, P.J., see Won, L., 6
Kurtz, H.J., see Skilling, S.R., 272

Laguzzi, R., see Merahi, N., 74
Lai, Y.Y. and Siegel, J.M., Corticotropin-releasing factor mediated muscle atonia in pons and medulla, 63
Larson, A.A., see Skilling, S.R., 272
Laruelle, M., Jaskiw, G.E., Lipska, B.K., Kolachana, B., Casanova, M.F., Kleinman, J.E. and Weinberger, D.R., D_1 and D_2 receptor modulation in rat striatum and nucleus accumbens after subchronic and chronic haloperidol treatment, 47
Lavie, V., Solomon, A., Ben-Bassat, S., Belkin, M. and Schwartz, M., Horseradish peroxidase labeling of growth cones and axons beyond the site of injury in injured rabbit optic nerve axons growing in their own environment, 1
Levant, B., see Myers, B., 325
Liebeskind, J.C., see Ben-Eliyahu, S., 304
Lipska, B.K., see Laruelle, M., 47
Liuzzi, F.J., see Smith, K.J., 79

Machado, A., see Vizuete, M.L., 231
Maisonneuve, I.M., Rossman, K.L., Keller Jr., R.W. and Glick, S.D., Acute and prolonged effects of ibogaine on brain dopamine metabolism and morphine-induced locomotor activity in rats, 69
Marek, P., see Ben-Eliyahu, S., 304
Markey, C.J., see Ikeda, H., 285
Markey, S.P., see Ikeda, H., 285
Markgraf, C.G., Green, E.J., Hurwitz, B.E., Morikawa, E., Dietrich, W.D., McCabe, P.M., Ginsberg, M.D. and Schneiderman, N., Sensorimotor and cognitive consequences of middle cerebral artery occlusion in rats, 238
Marrama, P., see Pich, E.M., 265
Martinez-Gomez, M., Chirino, R., Beyer, C., Komisaruk, B.R. and Pacheco, P., Visceral and postural reflexes evoked by genital stimulation in urethane-anesthetized female rats, 279
Masson-Pévet, M., see Gauer, F., 32
Masuda, Y., see Masuo, Y., 113
Masuo, Y., Ohtaki, T., Masuda, Y., Tsuda, M. and Fujino, M., Binding sites for pituitary adenyloate cyclase activating polypeptide (PACAP): comparison with vasoactive intestinal polypeptide (VIP) binding site localization in rat brain sections, 113
Matsunaga, K., see Matsuno, K., 315
Matsuno, K., Matsunaga, K. and Mita, S., Increase of extracellular acetylcholine level in rat frontal cortex induced by (+)*N* -allylnormetazochine as measured by brain microdialysis, 315
McAllen, R.M., Badoer, E., Shafton, A.D., Oldfield, B.J. and McKinley, M.J., Hemorrhage induces c-fos immunoreactivity in spinally projecting neurons of cat subretrofacial nucleus, 329
McCabe, P.M., see Markgraf, C.G., 238
McKinley, M.J., see McAllen, R.M., 329
Merahi, N., Orer, H.S. and Laguzzi, R., 5-HT_2 receptors in the nucleus tractus solitarius: characterisation and role in cardiovascular regulation in the rat, 74
Messori, B., see Pich, E.M., 265
Miller, R.F., see Henshel, D.S., 208
Mita, S., see Matsuno, K., 315
Mizisin, A.P., see Calcutt, N.A., 320
Mogil, J.S., see Ben-Eliyahu, S., 304
Mori, N. and Wada, J.A., Kindling of the massa intermedia of the thalamus in rats, 148
Morikawa, E., see Markgraf, C.G., 238
Muir, D., see Calcutt, N.A., 320
Myers, B., Levant, B., Bissette, G. and Nemeroff, C.B., Pharmacological specificity of the increase in neurotensin concentrations after antipsychotic drug treatment, 325

Nemeroff, C.B., see Myers, B., 325
Newman, G.C., Qi, H., Hospod, F.E. and Grundmann, K., Preservation of hippocampal brain slices with in vivo or in vitro hypothermia, 159
Newman, S.W., see Asmus, S.E., 199
Nojima, H., Kimura, I. and Kimura, M., Blocking action of succinylcholine with β-eudesmol on acetylcholine-activated channel activity at endplates of single muscle cells of adult mice, 337

Ohtaki, T., see Masuo, Y., 113
Okazaki, S., Sako, K. and Yonemasu, Y., Effects of intracarotid infusion of neuropeptide Y on cerebral blood flow in rats, 347
Oldfield, B.J., see McAllen, R.M., 329
Orer, H.S., see Merahi, N., 74

Pacheco, P., see Martinez-Gomez, M., 279
Pang, L. and Eyzaguirre, C., Different effects of hypoxia on the membrane potential and input resistance of isolated and clustered carotid body glomus cells, 167
Pévet, P., see Gauer, F., 32
Pich, E.M., Messori, B., Zoli, M., Ferraguti, F., Marrama, P., Biagini, G., Fuxe, K. and Agnati, L.F., Feeding and drinking responses to neuropeptide Y injections in the paraventricular hypothalamic nucleus of aged rats, 265
Pover, C.M., see Cameron, A.A., 151
Powell, H.C., see Calcutt, N.A., 320
Prasad, C., see Sakata, M., 309

Qi, H., see Newman, G.C., 159

Raggenbass, M., see Dubois-Dauphin, M., 124
Rao, Z.-r., see Chen, J., 333
Riekkinen Jr., P., Riekkinen, M., Sirviö, J. and Riekkinen, P., Effects of concurrent nicotinic antagonist and PCPA treatments on spatial and passive avoidance learning, 247
Riekkinen, M., see Riekkinen Jr., P., 247
Riekkinen, P., see Riekkinen Jr., P., 247
Robertson, D., see Cole, K.S., 223
Rogawski, M.A., see Coleman, M.H., 138
Rossman, K.L., see Maisonneuve, I.M., 69
Roth, R.H., see Al-Tikriti, M.S., 39

Sakata, M., Farooqui, S.M. and Prasad, C., Post-transcriptional regulation of loss of rat striatal D_2 dopamine receptor during aging, 309
Sako, K., see Okazaki, S., 347
Schneiderman, N., see Markgraf, C.G., 238
Schultz, E., see Jiao, S., 143
Schulz, C.A., see Herman, M.A.R., 257
Schwartz, M., see Lavie, V., 1
Seybold, V.S., see Jansen, E.M., 93
Shafton, A.D., see McAllen, R.M., 329
Shen, K.-F., see Crain, S.M., 13
Shi, J.-w., see Chen, J., 333
Siegel, J.M., see Lai, Y.Y., 63
Silver, I.A., see Deas, J., 251
Sirviö, J., see Riekkinen, Jr., P., 247
Skilling, S.R., Sun, X., Kurtz, H.J. and Larson, A.A., Selective potentiation of NMDA-induced activity and release of excitatory amino acids by dynorphin: possible roles in paralysis and neurotoxicity, 272
Smith, K.J., Kodama, R.T. and Liuzzi, F.J., Is CNS trauma a prerequisite for the elongation of CNS axons into denervated peripheral nerve?, 79
Smith, M.S., see Dohanics, J., 215
Solomon, A., see Lavie, V., 1
Steffen, V., see Vizuete, M.L., 231
Steinman, J.L., Carlton, S.M. and Willis, W.D., The segmental distribution of afferent fibers from the vaginal cervix and hypogastric nerve in rats, 25
Sternberg, W.F., see Ben-Eliyahu, S., 304
Sun, X., see Skilling, S.R., 272

Tamai, Y., see Kimura, A., 181
Tauc, L., see Baux, G., 155
Taylor, E.M., see Bernstein, I.L., 132
Thornhill, J., see Hugie, T., 57
Tokunaga, K. and Ishikawa, K., Evidence for different sites of monoamine oxidase and catechol-*O*-methyltransferase in the striatum, 164
Tribollet, E., see Dubois-Dauphin, M., 124
Tsuda, M., see Masuo, Y., 113

Vaccarino, A.L., see Ben-Eliyahu, S., 304
Verbalis, J.G., see Dohanics, J., 215
Vizuete, M.L., Steffen, V., Machado, A. and Cano, J., Effects of neonatal enucleation on catecholamine and serotonin turnover and amino acid levels in lateral geniculate nucleus and visual cortex of the adult rat, 231

Wada, J.A., see Mori, N., 148
Ward, S.J., see Jansen, E.M., 93
Weinberger, D.R., see Laruelle, M., 47
Widmer, H., see Dubois-Dauphin, M., 124

Willis, W.D., see Cameron, A.A., 151
Willis, W.D., see Steinman, J.L., 25
Wolff, J.A., see Jiao, S., 143
Won, L., Kontur, P.J., Choi, H.K., Hoffmann, P.C., Heller, B. and Heller, A., Acute and persistent effects of methamphetamine on developing monoaminergic neurons in reaggregate tissue culture, 6
Wu, M., see Kamada, T., 187

Yamaguchi, S.-i., see Coleman, M.H., 138
Yonemasu, Y., see Okazaki, S., 347

Zahm, D.S., An electron microscopic morphometric comparison of tyrosine hydroxylase immunoreactive innervation in the neostriatum and the nucleus accumbens core and shell, 341
Zoli, M., see Pich, E.M., 265

An electron microscopic morphometric comparison of tyrosine hydroxylase immunoreactive innervation in the neostriatum and the nucleus accumbens core and shell
D.S. Zahm (USA) 341

Effects of intracarotid infusion of neuropeptide Y on cerebral blood flow in rats
S. Okazaki, K. Sako and Y. Yonemasu (Japan) 347

Basic fibroblast growth factor protects striatal neurons in vitro from NMDA-receptor mediated excitotoxicity
A. Freese, S.P. Finklestein and M. DiFiglia (USA) 351

Restoration of diabetes-induced changes in enteric nerves by phorbol 12,13-dibutyrate
A. Belai and G. Burnstock (UK) 356

Author Index 359